

1. Kamata K., Kobayashi T. Changes in superoxide dismutase mRNA expression by streptozotocin-induced diabetes. *Br. J. Pharmacol.* 1996, 119: 583-589.
2. Kobayashi T., Kamata K. Relationships among cholesterol, superoxide anion and endothelium-dependent relaxation in diabetic rats. *Eur. J. Pharmacol.* 1999, 67: 213-222.
3. Kobayashi T., Kamata K. Effect of insulin treatment on the smooth muscle contractility and endothelium-dependent relaxation in rat aortae from established STZ-induced diabetes. *Br. J. Pharmacol.* 1999, 127: 835-842.
4. Kobayashi T., Matsumoto T., Kamata K. Mechanisms underlying the chronic pravastatin treatment-induced improvement in the impaired endothelium-dependent aortic relaxation seen in streptozotocin-induced diabetic rats. *Br. J. Pharmacol.*, 2000, 131: 231-238.
5. Kobayashi T., Kamata K. Effect of chronic insulin treatment on NO metabolism and endothelium-dependent relaxation in rats aortae from established STZ-induced diabetic rats. *Atherosclerosis.*, 2001, 155: 313-320.
6. Kobayashi T., Kamata K. Modulation by hydrogen peroxide of noradrenaline-induced contraction in aorta from streptozotocin-induced diabetic rat. *Eur. J. Pharmacol.*, 2002, 441: 83-89.
7. Shibutani M., Takahashi N., Kobayashi T., Uneyama C., Masutomi N., Nishikawa A., Hirose M. Molecular profiling of genes up-regulated during promotion by phenobarbital treatment in a medium-term rat liver bioassay. *Carcinogenesis.*, 2002, 23: 1047-1055.
8. Kobayashi T., Kamata K. Short-term insulin treatment and aortic expressions of IGF-1 receptor and VEGF mRNA in diabetic rats. *Am J Physiol Heart Circ Physiol.*, 2002, 283: H1761-1768.
9. Matsumoto T., Kobayashi T., Kamata K. Alterations in EDHF-type relaxation and phosphodiesterase activity in mesenteric arteries from diabetic rats. *Am J Physiol Heart Circ Physiol.*, 2003, 285: H283-291.
10. Kubota T., Kubota N., Moroi M., Terauchi Y., Kobayashi T., Kamata K., Suzuki R., Tobe K., Namiki A., Aizawa S., Nagai R., Kadowaki T., Yamaguchi T. Lack of insulin receptor substrate-2 causes progressive neointima formation in response to vessel injury. *Circulation.*, 2003, 107: 3073-3080.
11. Kanie N., Matsumoto T., Kobayashi T., Kamata K. Relationship between peroxisome proliferator-activated receptors α (PPAR α and PPAR γ) and endothelium-dependent relaxation in streptozotocin-induced diabetic rats. *Br J Pharmacol.* 2003, 140: 23-32.
12. Kobayashi T., Kaneda K., Kamata K. Possible involvement of IGF-1 receptor and IGF-binding protein in insulin-induced enhancement of noradrenaline response in diabetic rat aorta. *Br J Pharmacol.* 2003, 140: 285-294.
13. Matsumoto T., Wakabayashi K., Kobayashi T., Kamata K. Alterations in vascular endothelial function in the aorta and mesenteric artery in type II diabetic rats. *Can J Physiol Pharmacol.* 2004, 82(3):175-182.
14. Matsumoto T., Sato A., Suenaga H., Kobayashi T., Kamata K. Modulations of shear stress-induced contractile responses and agonist-induced vasodilation in hypercholesterolemic rats. *Atherosclerosis.*, 2004, 175: 31-38.
15. Kobayashi T., Matsumoto T., Ooishi K., Kamata K. Differential expressions of α

- 2D-adrenoceptor and eNOS in aortae from early and later stages of diabetes in Goto-Kakizaki rats. *Am J Physiol Heart Circ Physiol.*, 2004, 287: H135-143.
16. Matsumoto T., Oda S, Kobayashi T., Kamata K. Flow-Induced Endothelium-Dependent Vasoreactivity in Rat Mesenteric Arterial Bed. *J smooth muscle research.*, 2004, 40(1): 1-14.
 17. Matsumoto T., Wakabayashi K., Kobayashi T., Kamata K. Diabetes-related changes in cAMP-dependent protein kinase activity and decrease in relaxation response in rat mesenteric artery. *Am J Physiol Heart Circ Physiol.*, 2004, 287(3):H1064-1071.
 18. Matsumoto T., Yoshiyama S., Kobayashi T., Kamata K. Mechanisms underlying enhanced contractile response to endothelin-1 in diabetic rat basilar artery. *Peptides.* 2004, 25(11):1985-1994.
 19. Matsumoto T., Yoshiyama S., Wakabayashi K., Kobayashi T., Kamata K. Effects of chronic insulin on endothelial dysfunction of basilar arteries from established streptozotocin-diabetic rats. *Eur J Pharmacol.* 2004, 504(1-2):119-127.
 20. Matsumoto T., Yoshiyama S., Wakabayashi K., Kobayashi T., Kamata K. Effect of chronic insulin on cromakalim-induced relaxation in established streptozotocin-diabetic rat basilar artery. *Eur J Pharmacol.* 2004, 504(1-2):129-137.
 21. Kobayashi T., Taguchi K., Yasuhiro T., Matsumoto T., Kamata K. Impairment of PI3-K/Akt pathway underlies attenuated endothelial function in aorta of type 2 diabetic mouse model. *Hypertension.* 2004, 44(6):956-962.
 22. Kamata K, Kanie N, Matsumoto T, Kobayashi T. Endothelin-1-induced impairment of endothelium-dependent relaxation in aortae isolated from controls and diabetic rats. *J Cardiovasc Pharmacol.* 2004, 44, S186-S190.
 23. Kamata K., Kobayashi T., Matsumoto T., Kanie N., Oda S., Kaneda A., Sugiura M. Effects of Chronic Administration of Fruit Extract (Citrus unshiu MARC) on Endothelial Dysfunction in Streptozotocin-Induced Diabetic Rats. *Biol Pharm Bull.* 2005, 28(2):267-270.
 24. Matsumoto T, Kobayashi T, Kikuchi T, Honda T, Kamata K. Effects of dual-action genistein derivatives on relaxation in rat aorta. *J Smooth Muscle Res.* 2005 Feb;41(1):23-33.
 25. Endo K, Matsumoto T, Kobayashi T, Kasuya Y, Kamata K. Diabetes-related changes in contractile responses of stomach fundus to endothelin-1 in streptozotocin-induced diabetic rats. *J Smooth Muscle Res.* 2005 Feb;41(1):35-47.
 26. Matsumoto T, Wakabayashi K, Kobayashi T, Kamata K. Functional changes in adenylyl cyclases and associated decreases in relaxation responses in mesenteric arteries from diabetic rats. *Am J Physiol Heart Circ Physiol.* 2005 Nov;289(5):H2234-43.
 27. Matsumoto T, Kobayashi T, Wakabayashi K, Kamata K. Cilostazol improves endothelium-derived hyperpolarizing factor-type relaxation in mesenteric arteries from diabetic rats. *Am J Physiol Heart Circ Physiol.* 2005 Nov;289(5):H1933-40.
 28. Kobayashi T, Matsumoto T, Kamata K. IGF-1-induced enhancement of contractile response in organ-cultured aortae from diabetic rats is mediated by sustained thromboxane A₂ release from endothelial cells. *J Endocrinol,* 2005 Aug;186(2):367-76.
 29. Matsumoto T, Nishiyama M, Kobayashi T, Kasuya Y, Kamata K. Effect of phorbol

- 12,13-dibutyrate on smooth muscle tone in rat stomach fundus. *J Smooth Muscle Res.* 2005 Apr;41(2):107-116.
30. Kobayashi T, Oishi K, Hayashi Y, Matsumoto T, Kamata K. Changes in aortic endothelial gene expressions and relaxation responses following chronic short-term insulin treatment in diabetic rats *Atherosclerosis.*, 2006 Mar;185(1):47-57.
31. Kobayashi T, Hayashi Y, Taguchi K, Matsumoto T, Kamata K. Angiotensin II enhances contractile responses via PI3-kinase p110{delta} pathway in aortas from diabetic rats with systemic hyperinsulinemia. *Am J Physiol Heart Circ Physiol.* 2006; 291(2):H846-53.
32. Matsumoto T, Miyamori K, Kobayashi T, Kamata K. Specific impairment of endothelium-derived hyperpolarizing factor-type relaxation in mesenteric arteries from streptozotocin-induced diabetic mice. *Vascul Pharmacol.* 2006 Jun;44(6):450-60.
33. Matsumoto T, Kobayashi T, Kamata K. Mechanisms underlying the impaired EDHF-type relaxation response in mesenteric arteries from Otsuka Long-Evans Tokushima Fatty (OLETF) rats. *Eur J Pharmacol.* 2006 May 24;538(1-3):132-40.
34. Kamata K, Makino A, Kanie N, Oda S, Matsumoto T, Kobayashi T, Kikuchi T, Nishimura M, Honda T. Effects of anthocyanidin derivative (HK-008) on relaxation in rat perfused mesenteric bed. *J Smooth Muscle Res.* 2006 Jun;42(2-3):75-88.
35. Matsumoto T, Miyamori K, Kobayashi T, Kamata K. Apocynin normalizes hyperreactivity to phenylephrine in mesenteric arteries from cholesterol-fed mice by improving endothelium-derived hyperpolarizing factor response. *Free Radic Biol Med.* 2006 Oct 15;41(8):1289-303.
36. Matsumoto T, Kobayashi T, Kamata K. Mechanisms underlying lysophosphatidylcholine-induced potentiation of vascular contractions in the Otsuka Long-Evans Tokushima Fatty (OLETF) rat aorta. *Br J Pharmacol.* 2006 Dec;149(7):931-41.
37. Kamata K, Satoh T, Matsumoto T, Noguchi E, Taguchi K, Kobayashi T, Tanaka H, Shigenobu K. Enhancement of methoxamine-induced contractile responses of rat ventricular muscle in streptozotocin-induced diabetes is associated with alpha(1A) adrenoceptor upregulation. *Acta Physiol (Oxf).* 2006 Nov;188(3-4):173-83.
38. Kamata K, Hosokawa M, Matsumoto T, Kobayashi T. Acetylcholine-induced vasodilation in the perfused kidney of the streptozotocin-induced diabetic rat: role of prostacyclin. *J Smooth Muscle Res.* 2006 Oct;42(5):159-70.
39. Kamata K, Hosokawa M, Matsumoto T, Kobayashi T. Altered arachidonic acid-mediated responses in the perfused kidney of the streptozotocin-induced diabetic rat. *J Smooth Muscle Res.* 2006 Oct;42(5):171-87.
40. Matsumoto T, Kobayashi T, Wachi H, Seyama Y, Kamata K. Vascular NAD(P)H oxidase mediates endothelial dysfunction in basilar arteries from Otsuka Long-Evans Tokushima Fatty (OLETF) rats. *Atherosclerosis.* 2007 May;192(1):15-24.
41. Taguchi K, Kobayashi T, Hayashi Y, Matsumoto T, Kamata K. Enalapril improves impairment of SERCA-derived relaxation and enhancement of tyrosine nitration in diabetic rat aorta. *Eur J Pharmacol.* 2007 Feb 5;556(1-3):121-8.
42. Matsumoto T, Noguchi E, Kobayashi T, Kamata K. Mechanisms underlying the chronic pioglitazone treatment-induced improvement in the impaired endothelium-dependent

relaxation seen in aortas from diabetic rats. *Free Radic Biol Med.* 2007 Apr 1;42(7):993-1007.

43. Kobayashi T, Taguchi K, Takenouchi Y, Matsumoto T, Kamata K. Insulin-induced impairment via peroxynitrite production of endothelium-dependent relaxation and sarco/endoplasmic reticulum Ca²⁺-ATPase function in aortas from diabetic rats. *Free Radic Biol Med.* 2007 Aug 1;43(3):431-43.
44. Matsumoto T, Kakami M, Noguchi E, Kobayashi T, Kamata K. Imbalance between endothelium-derived relaxing and contracting factors in mesenteric arteries from aged OLETF rats, a model of type 2 diabetes. *Am J Physiol Heart Circ Physiol.* 2007 Sep;293(3):H1480-90.
45. Fukushima M, Hattori Y, Tsukada H, Koga K, Kajiwara E, Kawano K, Kobayashi T, Kamata K, Maitani Y. Adiponectin gene therapy of streptozotocin-induced diabetic mice using hydrodynamic injection. *J Gene Med.* 2007 Nov;9(11):976-85
46. Takenouchi Y, Kobayashi T, Matsumoto T, Kamata K. Possible involvement of Akt activity in endothelial dysfunction in type 2 diabetic mice. *J Pharmacol Sci.* 2008 Apr;106(4):600-8.
47. Matsumoto T, Kakami M, Kobayashi T, Kamata K. Gender differences in vascular reactivity to endothelin-1 (1-31) in mesenteric arteries from diabetic mice. *Peptides.* 2008 Aug;29(8):1338-46.
48. Matsumoto T, Noguchi E, Ishida K, Kobayashi T, Yamada N, Kamata K. Metformin normalizes endothelial function by suppressing vasoconstrictor prostanoids in mesenteric arteries from OLETF rats, a model of type 2 diabetes. *Am J Physiol Heart Circ Physiol.* 2008 Sep;295(3):H1165-H1176.
49. Kobayashi T, Nogami T, Taguchi K, Matsumoto T and Kamata K. Diabetic state, and high plasma insulin and Ang II combine to augment ET-1-induced contraction via ET_A-receptor and ERK. *Br J Pharmacol.* 2008 Dec;155(7):974-83
50. Matsumoto T, Noguchi E, Ishida K, Nakayama N, Kobayashi T, Kamata K. Cilostazol improves endothelial dysfunction by increasing endothelium-derived hyperpolarizing factor response in mesenteric arteries from Type 2 diabetic rats. *Eur J Pharmacol.* 2008 Dec 3;599(1-3):102-9
51. Kamata K, Ozawa Y, Kobayashi T, Matsumoto T. Effect of long-term streptozotocin-induced diabetes on coronary vasoconstriction in isolated perfused rat heart. *J Smooth Muscle Res.* 2008 Oct;44(5):177-88.
52. Matsumoto T, Takaoka E, Ishida K, Nakayama N, Noguchi E, Kobayashi T, Kamata K. Abnormalities of endothelium-dependent responses in mesenteric arteries from Otsuka Long-Evans Tokushima Fatty (OLETF) rats are improved by chronic treatment with thromboxane A₂ synthase inhibitor. *Atherosclerosis.* 2009 Jul;205(1):87-95.
53. Matsumoto T, Nakayama N, Ishida K, Kobayashi T, Kamata K. Eicosapentaenoic acid improves imbalance between vasodilator and vasoconstrictor actions of endothelium-derived factors in mesenteric arteries from rats at chronic stage of type 2 diabetes. *J Pharmacol Exp Ther.* 2009 Apr;329(1):324-34.
54. Matsumoto T, Ishida K, Nakayama N, Kobayashi T, Kamata K. Involvement of NO and MEK/ERK pathway in enhancement of endothelin-1-induced mesenteric artery contraction

- in later-stage type 2 diabetic Goto-Kakizaki rat. *Am J Physiol Heart Circ Physiol*. 2009 May;296(5):H1388-97.
55. Takenouchi Y, Kobayashi T, Matsumoto T, Kamata K. Gender differences in age-related endothelial function in the murine aorta. *Atherosclerosis*. 2009 Oct;206(2):397-404.
 56. Matsumoto T, Ishida K, Kobayashi T, Kamata K. Pyrrolidine dithiocarbamate reduces vascular prostanoid-induced responses in aged type 2 diabetic rat model. *J Pharmacol Sci*. 2009 Jul;110(3):326-33.
 57. Kamata K, Ozawa Y, Kobayashi T, Matsumoto T. Effect of N-epsilon-(carboxymethyl)lysine on coronary vasoconstriction in isolated perfused hearts from control and streptozotocin-induced diabetic rats. *J Smooth Muscle Res*. 2009 Jun;45(2-3):125-37.
 58. Matsumoto T, Ishida K, Taguchi K, Kobayashi T, Kamata K. Mechanisms underlying enhanced vasorelaxant response to protease-activated receptor 2-activating peptide in type 2 diabetic Goto-Kakizaki rat mesenteric artery. *Peptides*. 2009 Sep;30(9):1729-34..
 59. Kobayashi T, Taguchi K, Nemoto S, Nogami T, Matsumoto T, Kamata K. Activation of the PDK1/Akt /eNOS pathway involved in aortic endothelial function differs between hyperinsulinemic and insulin-deficient diabetic rats. *Am J Physiol Heart Circ Physiol*. 2009 Nov;297(5):H1767-75.
 60. Takenouchi Y, Kobayashi T, Taguchi K, Matsumoto T, Kamata K. Gender Differences in Endothelial Function in Aortas From Type 2 Diabetic Model Mice. *J Pharmacol Sci*. 2009 Sep;111(1):91-9.
 61. Matsumoto T, Ozawa Y, Taguchi K, Kobayashi T, Kamata K. Chronic treatment with losartan (angiotensin II type 1 receptor antagonist) normalizes enhanced acetylcholine-induced coronary vasoconstriction in isolated perfused hearts of type 2 diabetic OLETF rats. *J Smooth Muscle Res*. 2009 Oct;45(5):197-208
 62. Matsumoto T, Ozawa Y, Taguchi K, Kobayashi T, Kamata K. Diabetes-associated changes and role of N(varepsilon)-(carboxymethyl)lysine in big ET-1-induced coronary vasoconstriction. *Peptides*. 2010 Feb;31(2):346-53.
 63. Matsumoto T, Ishida K, Taguchi K, Kobayashi T, Kamata K. Short-term angiotensin-1 receptor antagonism in type 2 diabetic Goto-Kakizaki rats normalizes endothelin-1-induced mesenteric artery contraction. *Peptides*. 2010 Apr;31(4):609-17.
 64. Matsumoto T, Ishida K, Taguchi K, Kobayashi T, Kamata K. Losartan Normalizes Endothelium-Derived Hyperpolarizing Factor-Mediated Relaxation by Activating Ca(2+)-Activated K(+) Channels in Mesenteric Artery From Type 2 Diabetic GK Rat. *J Pharmacol Sci*. 2010 Mar 19;112(3):299-309.
 65. Matsumoto T, Ishida K, Nakayama N, Taguchi K, Kobayashi T, Kamata K. Mechanisms underlying the losartan treatment-induced improvement in the endothelial dysfunction seen in mesenteric arteries from type 2 diabetic rats. *Pharmacol Res*. 2010 Sep;62(3):271-81..
 66. Takenouchi Y, Kobayashi T, Taguchi K, Matsumoto T, Kamata K. Relationship among superoxide-related enzyme, PPARs, and endothelium-dependent relaxation in murine aortas previously organ-cultured in high-glucose conditions. *Can J Physiol Pharmacol*. 2010 Jul;88(7):760-9.
 67. Matsumoto T, Kobayashi T, Ishida K, Hirasawa Y, Morita H, Honda T, Kamata K.

- Vasodilator effect of Cassiarin A, a novel antiplasmodial alkaloid from *Cassia siamea*, in rat isolated mesenteric artery. *Biol Pharm Bull.* 2010;33(5):844-8.
68. Matsumoto T, Kobayashi T, Ishida K, Taguchi K, Kamata K. Enhancement of mesenteric artery contraction to 5-HT depends on Rho kinase and Src kinase pathways in the ob/ob mouse model of type 2 diabetes. *Br J Pharmacol.* 2010 Jul;160(5):1092-104.
 69. Sasaki Y, Suzuki M, Matsumoto T, Hosokawa T, Kobayashi T, Kamata K, Nagumo S. Vasorelaxant activity of Sappan Lignum constituents and extracts on rat aorta and mesenteric artery. *Biol Pharm Bull.* 2010;33(9):1555-60.
 70. Takenouchi Y, Kobayashi T, Taguchi K, Matsumoto T, Kamata K. Gender differences in vascular reactivity of aortas from streptozotocin-induced diabetic mice. *Biol Pharm Bull.* 2010;33(10):1692-7.
 71. Taguchi K, Kobayashi T, Matsumoto T, Kamata K. Dysfunction of endothelium-dependent relaxation to insulin via PKC-mediated GRK2/Akt activation in aortas of ob/ob mice. *Am J Physiol Heart Circ Physiol.* 2011 Aug;301(2):H571-83.
 72. Kubota T, Kubota N, Kumagai H, Yamaguchi S, Kozono H, Takahashi T, Inoue M, Itoh S, Takamoto I, Sasako T, Kumagai K, Kawai T, Hashimoto S, Kobayashi T, Sato M, Tokuyama K, Nishimura S, Tsunoda M, Ide T, Murakami K, Yamazaki T, Ezaki O, Kawamura K, Masuda H, Moroi M, Sugi K, Oike Y, Shimokawa H, Yanagihara N, Tsutsui M, Terauchi Y, Tobe K, Nagai R, Kamata K, Inoue K, Kodama T, Ueki K, Kadowaki T. Impaired insulin signaling in endothelial cells reduces insulin-induced glucose uptake by skeletal muscle. *Cell Metab.* 2011 Mar 2;13(3):294-307.
 73. Taguchi K, Kobayashi T, Takenouchi Y, Matsumoto T, Kamata K. Angiotensin II causes endothelial dysfunction via the GRK2/Akt/eNOS pathway in aortas from a murine type 2 diabetic model. *Pharmacol Res.* 2011 Nov;64(5):535-46.
 74. Ishida K, Matsumoto T, Taguchi K, Kamata K, Kobayashi T. Mechanisms underlying altered extracellular nucleotide-induced contractions in mesenteric arteries from rats in later-stage type 2 diabetes: effect of ANG II type 1 receptor antagonism. *Am J Physiol Heart Circ Physiol.* 2011 Nov;301(5):H1850-61.
 75. Nemoto S, Kobayashi T, Taguchi K, Matsumoto T, Kamata K. Losartan improves aortic endothelium-dependent relaxation via proline-rich tyrosine kinase 2/Src/Akt pathway in type 2 diabetic Goto-Kakizaki rats. *Am J Physiol Heart Circ Physiol.* 2011 Dec;301(6):H2383-94.
 76. Taguchi K, Matsumoto T, Kamata K, Kobayashi T. Akt/eNOS pathway activation in endothelium-dependent relaxation is preserved in aortas from female, but not from male, type 2 diabetic mice. *Pharmacol Res.* 2012 Jan;65(1):56-65.
 77. Ishida K, Matsumoto T, Taguchi K, Kamata K, Kobayashi T. Pravastatin normalizes EDCF-mediated response via suppression of Rho-kinase signalling in mesenteric artery from aged type 2 diabetic rat. *Acta Physiol (Oxf).* 2012 Jun;205(2):255-65.
 78. Ishida K, Matsumoto T, Taguchi K, Kamata K, Kobayashi T. Protein kinase C delta contributes to increase in EP3 agonist-induced contraction in mesenteric arteries from type 2 diabetic Goto-Kakizaki rats. *Pflugers Arch.* 2012 Apr;463(4):593-602.
 79. Taguchi K, Matsumoto T, Kamata K, Kobayashi T. Angiotensin II type 2 receptor-dependent increase in nitric oxide synthase activity in the endothelium of db/db

- mice is mediated via a MEK pathway. *Pharmacol Res.* 2012 Jul;66(1):41-50.
80. Nemoto S, Taguchi K, Matsumoto T, Kamata K and Kobayashi T. Aminoguanidine normalizes ET-1-induced aortic contraction in type 2 diabetic Otsuka Long-Evans Tokushima Fatty (OLETF) rats by suppressing Jab1-mediated increase in ET_A-receptor expression. *Peptides.* 2012 Jan;33(1):109-19.
 81. Taguchi K, Matsumoto T, Kamata K, Kobayashi T. G Protein-Coupled Receptor Kinase 2, With β -Arrestin 2, Impairs Insulin-Induced Akt/Endothelial Nitric Oxide Synthase Signaling in ob/ob Mouse Aorta. *Diabetes.* 2012 Aug;61(8):1978-85.
 82. Taguchi K, Matsumoto T, Kamata K, Kobayashi T. Inhibitor of G Protein-Coupled Receptor Kinase 2 Normalizes Vascular Endothelial Function in Type 2 Diabetic Mice by Improving β -Arrestin 2 Translocation and Ameliorating Akt/eNOS Signal Dysfunction. *Endocrinology.* 2012 Jul;153(7):2985-96.
 83. Kobayashi T, Nemoto S, Ishida K, Taguchi K, Matsumoto T, Kamata K. Involvement of CaM kinase II in the impairment of endothelial function and eNOS activity in aortas of Type 2 diabetic rats. *Clin Sci (Lond).* 2012 Sep 1;123(6):375-86.
 84. Taguchi K, Morishige A, Matsumoto T, Kamata K, Kobayashi T. Enhanced estradiol-induced vasorelaxation in aortas from type 2 diabetic mice may reflect a compensatory role of p38 MAPK-mediated eNOS activation. *Pflugers Arch.* 2012 Aug;464(2):205-15.
 85. Nemoto S, Taguchi K, Matsumoto T, Kamata K, Kobayashi T. Pravastatin normalizes ET-1-induced contraction in aorta of type 2 diabetic OLETF rat by suppressing kinase suppressor of Ras (KSR)-1/ERK complex. *Am J Physiol Heart Circ Physiol.* 2012 Oct;303(7):H893-902.
 86. Taguchi K, Matsumoto T, Kamata K, Kobayashi T. Suppressed G-protein-coupled receptor kinase 2 activity protects female diabetic-mouse aorta against endothelial dysfunction. *Acta Physiol (Oxf).* 2013 Jan;207(1):142-55.
 87. Ishida K, Matsumoto T, Taguchi K, Kamata K, Kobayashi T. Mechanisms underlying reduced P2Y(1)-receptor-mediated relaxation in superior mesenteric arteries from long-term streptozotocin-induced diabetic rats. *Acta Physiol (Oxf).* 2013 Jan;207(1):130-41.
 88. Matsumoto T, Watanabe S, Kawamura R, Taguchi K, Kobayashi T. Enhanced uridine adenosine tetraphosphate-induced contraction in renal artery from type 2 diabetic Goto-Kakizaki rats due to activated cyclooxygenase/thromboxane receptor axis. *Pflugers Arch.* 2013 Jul 31. [Epub ahead of print] PubMed PMID: 23900807.
 89. Tagashira H, Matsumoto T, Taguchi K, Zhang C, Han F, Ishida K, Nemoto S, Kobayashi T, Fukunaga K. Vascular Endothelial α (1)-Receptor Stimulation With SA4503 Rescues Aortic Relaxation via Akt/eNOS Signaling in Ovariectomized Rats With Aortic Banding. *Circ J.* 2013 Aug 20. [Epub ahead of print] PubMed PMID:23965801.